

Table 1: Comparison of characteristics found in the group with PNEE remission (event freedom) to group with continued events using data collected at initial clinic visit, one-month follow-up, three-month follow-up, and 12-month follow-up. Variable rates of patient participation at one-month and three-months necessitates adjusting denominator (representing participation in that question) to display accurate percentages.

Table 1	12 Month Remission (n=16)		12 Month Combined Remission and Improvement (n=50)		12 Month No Change and Worsened PNEE (n=7)		12 Month Acceptance (n=42)		Summed Total*		
	N=	Percent of Subset	N=	Percent of Subset	N=	Percent of Subset	N=	Percent of subset	Positive Responses	Total Responses to Question	Percent
Acceptance of Diagnosis at One-Month	9	56	23	46	1	14	22	52	24	45	53
Acceptance of Diagnosis at Three-Months	8	50	29	58	3	43	26	62	32	42	76
Acceptance of Diagnosis at 12-Months	13	81	38	76	3	43	42	100	42	57	74
Linked with Counseling at Clinic Visit	6	38	16	32	0	0	10	24	16	56	29
Linked with Counseling at One-Month	12	75	28	56	3	43	23	55	31	46	67
Linked with Counseling at Three-Months	8	50	25	50	3	43	21	50	28	42	67
Linked with Counseling at 12-Months	12	75	39	78	3	43	33	79	41	57	72
Documented on EEG	7	44	21	42	4	57	17	40	25	57	44
Remission at One-Month									11	44	25
Remission at Three-Months									19	42	45
Remission at 12-Months									16	57	28
Improved and Remission at One-Month									32	44	73
Improved and Remission at Three-Months									36	42	86
Improved and Remission at 12-Months									50	57	88
ED Utilization at 12-Months	0	0	5	10	3	43			8	57	14
Hospitalizations at 12-Months	0	0	2	4	0	0			1	57	2

*there was variability in responses to individual follow-up questions