

Table 2. Incidence of on-treatment suicidality reports in pediatric patients (aged 6 to <12 years) based on the C-SSRS during the Study 311 Core Study and Extension A (Safety Analysis Set^a)

	POS (n=122)	SGS (n=43)	PGTCS (n=27)	Total (N=149)
Patients with ≥ 1 positive ideation, n (%)	2 (1.6)	0 (0.0)	2 (7.4)	4 (2.7)
Wish to be dead	2 (1.6)	0 (0.0)	1 (3.7)	3 (2.0)
Non-specific active thoughts	2 (1.6)	0 (0.0)	2 (7.4)	4 (2.7)
Active thoughts without intent to act	2 (1.6)	0 (0.0)	1 (3.7)	3 (2.0)
Active thoughts with some intent – no plan	1 (0.8)	0 (0.0)	0 (0.0)	1 (0.7)
Active thoughts with plan and intent	1 (0.8)	0 (0.0)	0 (0.0)	1 (0.7)
Patients with ≥ 1 positive behavior, n (%)	1 (0.8)	0 (0.0)	0 (0.0)	1 (0.7)
Preparatory acts or behavior	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Aborted attempt	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Interrupted attempt	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Actual attempt	1 (0.8) ^b	0 (0.0)	0 (0.0)	1 (0.7) ^b
Completed suicide	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Suicidality, n (%)	2 (1.6)	0 (0.0)	2 (7.4)	4 (2.7)

Patients were counted in each applicable category. An on-treatment report of suicidal ideation, suicidal behavior, or suicidality was an answer of “Yes” to any question in the respective category of the C-SSRS during perampanel treatment

^aIncludes patients with ≥ 1 on-treatment assessment of the C-SSRS

^bIt was confirmed that the C-SSRS had been erroneously reported by this patient, and as such, the investigator judged that there was no suicidal risk

C-SSRS, Columbia-Suicide Severity Rating Scale; PGTCS, primary generalized tonic-clonic seizures; POS, partial-onset seizures; SGS, secondarily generalized seizures