

Table 1. Efficacy outcomes according to baseline SSF quartile

	Arm 1					Arm 2				
	All	Q1	Q2	Q3	Q4	All	Q1	Q2	Q3	Q4
Retention rate, n (%)	n = 44 36 (81.8)	n = 16 13 (81.3)	n = 8 7 (87.5)	n = 9 7 (77.8)	n = 11 9 (81.8)	n = 58 37 (63.8)	n = 10 6 (60.0)	n = 17 12 (70.6)	n = 16 10 (62.5)	n = 14 9 (64.3)
Change ^a in SSF, median %	n = 40 -72.8	n = 15 -84.8	n = 7 -40.9	n = 7 -74.0	n = 11 -59.2	n = 49 -22.8	n = 9 82.9	n = 14 -45.7	n = 14 -13.7	n = 12 -42.4
50% responder rate, ^b n (%)	n = 40 25 (62.5)	n = 15 11 (73.3)	n = 7 3 (42.9)	n = 7 4 (57.1)	n = 11 7 (63.6)	n = 52 20 (38.5)	n = 10 4 (40.0)	n = 14 6 (42.9)	n = 15 4 (26.7)	n = 12 6 (50.0)
Seizure freedom rate, n (%)	n = 41 10 (24.4)	n = 15 6 (40.0)	n = 7 0	n = 8 1 (12.5)	n = 11 3 (27.3)	n = 55 5 (9.1)	n = 10 2 (20.0)	n = 16 2 (12.5)	n = 15 1 (6.7)	n = 13 0

Q1 to Q4 = lowest to highest quartiles of seizure frequency.

Median baseline SSF in Arm 1: Q1, 0.86; Q2, 1.60; Q3, 3.98; Q4, 12.48.

Median baseline SSF in Arm 2: Q1, 0.80; Q2, 1.84; Q3, 3.92; Q4, 15.56.

^aChange from baseline.

^bProportion of patients with a \geq 50% reduction from baseline in SSF.

Q1–4: quartile 1–4; SSF: standardized seizure frequency (seizures per 4 weeks).