

**Table 2.** Approach to epilepsy management in TSC children according to geographical distribution- time of treatment initiation

Type of the survey question is indicated in the square brackets: single-choice or open (free text). One response from African clinician who does not take care of TSC children below 2 years of age was excluded from this analysis.

In case of a newborn/infant with TSC when would you introduce AEDs? (n=60) [single choice]	After clinical seizures	When EEG shows epileptic discharges but before clinical seizures	Other
		20 / 60 33.3%	31 / 60 51.7%
<b>Geographical distribution</b>			
<b>Europe</b> among European respondents (n=34)	10 / 34 (29.4%)	<b>21 / 34</b> <b>(61.8%)</b>	3 / 34 (8.8%)
<b>Asia</b> among Asian respondents (n=10)	<b>5 / 10</b> <b>(50.0%)</b>	4 / 10 (40.0%)	1 / 10 (10.0%)
<b>North America</b> among North American respondents (n=12)	4 / 12 (33.3%)	3 / 12 (25.0%)	<b>5 / 12</b> <b>(41.7%)</b>
<b>Australia</b> among Australian respondents (n=2)	-	<b>2 / 2</b> <b>(100%)</b>	-
<b>South America</b> among South American respondents (n=2)	<b>1 / 2</b> <b>(50.0%)</b>	<b>1 / 2</b> <b>(50.0%)</b>	-