

Table 1. Overview of 50% responder rates, seizure-freedom rates and percentage reduction in seizure frequency per 28 days in patients with POS by baseline concomitant AED and perampanel maintenance dose (Full Analysis Set).

	Levetiracetam (n=34)					Carbamazepine (n=20)					Oxcarbazepine (n=17)					Lamotrigine (n=7)					Valproic acid (n=4)				
	Perampanel mg/day																								
	4 n=20	6 n=11	8 n=3	10 n=0	12 n=0	4 n=7	6 n=6	8 n=6	10 n=1	12 n=0	4 n=7	6 n=7	8 n=1	10 n=1	12 n=1	4 n=6	6 n=0	8 n=1	10 n=0	12 n=0	4 n=2	6 n=2	8 n=0	10 n=0	12 n=0
50% responder, % , n	85.0 17/20	81.8 9/11	66.7 2/3	NA		100.0 7/7	83.3 5/6	50.0 3/6	0 0	NA	100.0 7/7	100.0 7/7	0 0	0 0	0 0	100.0 6/6	NA	0 0			100.0 2/2	50.0 1/2			
Seizure-freedom rate, % , n	50.0 10/20	36.4 4/11	0 0			71.4 5/7	50.0 3/6	16.7 1/6	0 0		85.7 6/7	71.4 5/7	0 0	0 0	0 0	50.0 3/6		0 0			100.0 2/2	0 0			
Median percent reduction in seizure freq. per 28 days, % , n	100.0 19/20	83.1 11/11	75.0 3/3			100.0 7/7	95.9 6/6	40.7 6/6	37.4 1/1		100.0 7/7	100.0 7/7	16.7 1/1	-63.3 1/1	14.3 1/1	96.9 6/6		-1457.3 1/1			100.0 2/2	57.3 2/2			

NA: not applicable, no patients in this group received the stated AED. AED, antiepileptic drug; freq. frequency; POS, partial-onset seizures