

Table 1. Baseline characteristics of the study patients (safety analysis set)

Patient background factors	Category	Adults (n=267)	%	Children (n=98)	%	Overall (n=365)	%
Gender	Male	124	46.44	46	46.94	170	46.58
	Female ^{a)}	143	53.56	51	52.04	194	53.15
	Unknown	0	0.00	1	1.02	1	0.27
Age (years)	Mean ± SD	36.36±20.10		11.38±3.59		29.65±20.53	
Weight (kg)	Mean ± SD	55.97±13.87		37.48±16.13		49.63±17.09	
	Unknown	100	37.45	11	11.22	111	30.41
Age at the onset of epilepsy (years)	Mean ± SD	20.26±20.37		7.60±4.64		16.65±18.31	
	Unknown	24	8.99	1	1.02	25	6.85
Duration of epilepsy (years)	Mean ± SD	15.03±13.62		3.70±3.71		11.80±12.75	
	Unknown	24	8.99	1	1.02	25	6.85
Types of seizures ^{b, c)}	Primary Tonic-clonic seizures	267	100.00	98	100.00	365	100.00
	Myoclonic seizures	51	19.10	25	25.51	76	20.82
	Complex partial seizures	15	5.62	11	11.22	26	7.12
	Absence seizures	19	7.12	7	7.14	26	7.12
	Secondary generalized seizures	14	5.24	7	7.14	21	5.75
	Simple partial seizures	9	3.37	2	2.04	11	3.01
Baseline seizure frequency of Tonic-clonic seizures	< 1 per 4 weeks	103	38.58	31	31.63	134	36.71
	≥ 1 per 4 weeks	156	58.43	65	66.33	221	60.55
	Unknown	8	3.00	2	2.04	10	2.74
Mean daily dose of levetiracetam (mg)	Mean ± SD	837.76±414.86		662.75±377.51		790.77±412.05	
Dosage form	Tablet	229	85.77	55	56.12	284	77.81
	Dry syrup	33	12.36	40	40.82	73	20.00
	Tablet and dry syrup	5	1.87	3	3.06	8	2.19
Number of concomitant antiepileptics at the start of treatment	1	184	68.91	78	79.59	262	71.78
	2	57	21.35	12	12.24	69	18.90
	3	18	6.74	6	6.12	24	6.58
	4	6	2.25	2	2.04	8	2.19
	5 or more	2	0.75	0	0.00	2	0.55
Concomitant antiepileptic drugs at the start of treatment ^{b, d)}	Sodium valproate	173	64.79	71	72.45	244	66.85
	Carbamazepine	38	14.23	9	9.18	47	12.88
	Lamotrigine	29	10.86	10	10.20	39	10.68
	Clonazepam	30	11.24	6	6.12	36	9.86
	Zonisamide	24	8.99	6	6.12	30	8.22
	Phenobarbital	19	7.12	7	7.14	26	7.12
	Clobazam	15	5.62	9	9.18	24	6.58
	Phenytoin	23	8.61	0	0.00	23	6.30
	Lacosamide	8	3.00	2	2.04	10	2.74

a) Including 2 pregnant and 2 lactating adult females (there is 1 overlapping)

b) There is some overlapping

c) Seizure types reported for 10 or more patients in total

d) Antiepileptic drugs that were used by 10 or more patients in total