

Table. Anxiety and depression prevalence estimates & 95% Confidence Intervals by instrument and epilepsy status

		Diagnosis of Epilepsy		Seizure Free in Past 6 Months*			
Instrument (positive values)	Overall (N = 422)	No/Uncertain (N = 58)	Yes (N = 364)	Uncertain (N = 15)	Yes (N = 188)	No (N = 161)	P Value**
GAD-7 (≥ 10)	0.24 0.21-0.29	0.22 0.14-0.35	0.25 0.21-0.30	0.40 0.20-0.64	0.17 0.12-0.23	0.32 0.26-0.40	0.0011
GAD-2 (> 2)	0.31 0.27-0.35	0.35 0.24-0.47	0.30 0.26-0.35	0.47 0.25-0.70	0.20 0.15-0.27	0.40 0.33-0.48	<0.0001
GAD-SI (≥ 2)	0.23 0.19-0.27	0.31 0.21-0.44	0.21 0.18-0.26	0.33 0.15-0.58	0.18 0.13-0.24	0.25 0.19-0.32	0.11
NDDI-E (≥ 16)	0.20 0.17-0.25	0.26 0.16-0.38	0.20 0.16-0.24	0.33 0.15-0.58	0.14 0.096-0.20	0.25 0.19-0.32	0.0094
NDDIE-2 (> 4)	0.20 0.16-0.24	0.28 0.18-0.40	0.18 0.15-0.23	0.40 0.20-0.64	0.11 0.070-0.16	0.26 0.19-0.33	0.0004

Confidence interval, CI; Generalized Anxiety Disorder-7, GAD-7; Generalized Anxiety Disorder-2, GAD-2; Single-item GAD, GAD-SI; Neurological Disorders Depression Inventory-Epilepsy, NDDI-E; Neurological Disorders Depression Inventory Epilepsy-2, NDDIE-2

*Among those with diagnosis of epilepsy

**P values based on Fisher exact test for comparison of seizure free and non-seizure free subgroups