

Table 2: Impact on child nutritional status among children under 2 years in control and intervention groups by DID estimates

	Intervention		Control		Unadjusted Difference in difference in % (95% CIs) ^a	P-value	Adjusted Difference in difference in % (95% CIs) ^{a, b}	P-value
	Baseline	Endline	Baseline	Endline				
Stunting	418 (28.6)	383 (24.2)	327 (22.3)	372 (23.5)	-5.7 (-10, -1.3)	0.010	-5.0 (-9.9, -0.2)	0.041
n	1461	1584	1467	1582				
Wasting	121 (8.4)	82 (5.2)	207 (14.2)	190 (12.1)	-1.1 (-4.1, 1.9)	0.455	-1.7 (-5.1, 1.6)	0.315
n	1447	1592	1458	1567				
Underweight	217 (14.9)	211 (13.2)	222 (15.1)	309 (19.3)	-5.8 (-9.5, -2.2)	0.002	-4.6 (-8.6, -0.5)	0.028
n	1461	1596	1467	1598				

a. Unadjusted and adjusted differences in difference were obtained from mixed linear regression models with an interaction term between health area (intervention vs control) and time (endline vs baseline).

b. Multivariable models adjusted for child sex, age, maternal age, education, skilled birth attendant, facility births, IYCF practices, maternal BMI, and household characteristics.

DiD, difference-in-differences.