

Table 2: Pulmonary Hypertension Outcomes by Bronchopulmonary Dysplasia Definition

Measure	Moderate (n=105)	Severe (n=292)	Grade 1 (n=246)	Grade 2 (n=98)	Grade 3 (n=53)	P value*
BPD-PH during hospitalization, n (%)	20 (19)	119 (41)	64 (26)	44 (45)	31 (58)	0.02
BPD-PH persistence at discharge, n (%)	7(7)	70 (24)	35 (14)	22 (22)	20 (38)	0.04
BPD-PH associated mortality, n (%)	0 (0)	25 (9)	6 (2)	7 (7)	12 (23)	0.002
BPD-PH or death, n (%)	20 (19)	126 (43)	64 (26)	45 (46)	37 (70)	<0.001
BPD-PH Pharmacology, n (%)	12 (11)	115 (39)	45 (18)	42 (43)	40 (76)	<0.001
Sildenafil, n (%)	7 (7)	72 (25)	29 (12)	30 (31)	20 (38)	0.048
Inhaled nitric oxide, n (%)	7 (7)	101 (35)	34 (14)	36 (37)	38 (72)	<0.001
Bosentan, n (%)	0 (0)	14 (5)	3 (1)	6 (6)	5 (9)	0.17

*p value by chi square analysis between severe BPD and grade 3 BPD