**Table 1:** Demographic and clinical characteristics of adolescents and young adults treated in the emergency department for cannabis hyperemesis syndrome

Demographic and Clinical	
Characteristics	n=4,571
Age categories	
13-18 years	3,684 (80.6%)
19-21 years	887 (19.4%)
Age (Years)	17.0 [16.0, 18.0]
Sex	
Female	2,888 (63.2%)
Male	1,682 (36.8%)
Race and Ethnicity	
Non-Hispanic White	1,769 (38.7%)
Non-Hispanic Black	1,132 (24.8%)
Hispanic	1,298 (28.4%)
Other	372 (8.1%)
Insurance Status	
Public	2,878 (63.0%)
Private	1,338 (29.3%)
Other	355 (7.8%)
Patient Urbanicity (based on zip code)	
Not Urban	266 (5.9%)
Urban/Suburban	4,264 (94.1%)
Overall Child Opportunity Level	
Very Low	1,433 (31.4%)
Low	935 (20.5%)
Moderate	789 (17.3%)
High	774 (17.0%)
Very High	634 (13.9%)
Disposition	
Discharged	2,666 (58.3%)
Floor/Observation unit	1,878 (41.1%)
ICU	27 (0.6%)
ICD-10 Diagnoses <sup>1</sup>	
Cannabis abuse	1,124 (24.6%)
Cannabis dependence	183 (4.0%)
Cannabis poisoning <sup>2</sup>	148 (3.2%)
Cannabis use, unspecified	3,159 (69.1%)

Values in table represent frequency (percent) or median [interquartile range]

<sup>&</sup>lt;sup>1</sup> Not mutually exclusive

<sup>&</sup>lt;sup>2</sup> Poisoning by, adverse effect of and underdosing of cannabis (derivatives)