

**Table 2: Clinical actions by CDC Congenital Syphilis Category**

Intervention	CDC Risk Category				Total
	Unlikely	Less Likely	Possible	Highly Probable /Proven	
RPR test					
Reactive	6 (13%)	31 (41%)	23 (48%)	9 (90%)	69 (38.5%)
Non-Reactive	25 (53%)	38 (51%)	25 (52%)	1 (10%)	89 (49.7%)
RPR not done	16 (34%)	5 (7%)	0 (0.0%)	0 (0.0%)	21 (11.7%)
Clinical workup					
Complete *	1 (2%)	3 (4%)	19 (40%)	7 (70%)	30 (16.8%)
Partial**	10 (21%)	21 (28%)	20 (42%)	3 (30%)	54 (30.2%)
None	36 (77%)	50 (68%)	9 (19%)	0 (0.0%)	95 (53.1%)
Penicillin treatment					
10 days	0 (0.0%)	5 (7%)	28 (58%)	10 (100%)	43 (24.0%)
1 dose	6 (13%)	32 (43%)	13 (27%)	0 (0.0%)	51 (28.5%)
None	41(87%)	37 (50%)	7 (15%)	0 (0.0%)	85 (47.5%)
Total	47(26%)	74 (41%)	48 (27%)	10 (6%)	179 (100.0%)

\* Complete blood count, long bone x-rays and lumbar puncture performed.

\*\* 1 or 2 of preceding tests resulted.