

Overall Likert Confidence Scores Pre- and Post-Intervention Using Wicxon Signed-Rank Test	Pre-Intervention Confident, n(%)	Post-Intervention Confident, n(%)	p-value
I can effectively screen for substance use disorder, including alcohol and opioids.	21 (20.8%)	34 (50%)	0.008*
I can recognize the symptoms of opioid withdrawal.	25 (24.8%)	36 (53.7%)	0.01*
I can choose the most appropriate medication to treat opioid use disorder using my patient's age.	25 (25%)	36 (53.7%)	<0.001*
I can effectively communicate with families of my patients while maintaining their confidentiality	40 (39.6%)	35 (52.2%)	0.001*
I feel comfortable providing counseling to patients and/or families about medications for treatment of opioid use disorder including long-term monitoring and risk/benefits.	28 (27.7%)	31 (45.6%)	0.003*

Overall Knowledge Scores Pre- and Post-Intervention Using McNemar Test	Pre-Intervention Correct, n(%)	Post-Intervention Correct, n(%)	p-value
Which of the following is the correct scoring device for opioid withdrawal symptoms?	70 (68.6%)	66 (98.5%)	0.021*
The opioid withdrawal scoring tool includes which of the following?	53 (52.4%)	47 (70.1%)	0.289
Which of the following is the best treatment for a 16-year-old with opioid use disorder?	67 (67%)	54 (83.1%)	<0.001*
True or false: In Kentucky, confidentiality laws enforce disclosure of treatment for substance use disorder to a guardian.	75 (73.5%)	47 (70.1%)	1