

<b>Didactic Likert Scale Confidence Pre- and Post Intervention Using Wilcoxon Signed-Rank Test</b>	<b>Pre-Intervention Confident, n(%)</b>	<b>Post-Intervention Confident, n(%)</b>	<b>p-value</b>
I can effectively screen for substance use disorder, including alcohol and opioids	12 (22.2%)	17 (53.1%)	0.033*
I can recognize the symptoms of opioid withdrawal	14 (25.9%)	16 (51.6%)	0.162
I can choose the most appropriate medication to treat opioid use disorder using my patient's age.	12 (22.7%)	17 (53.1%)	0.013*
I can effectively communicate with families of my patients while maintaining their confidentiality.	23 (42.6%)	19 (61.3%)	0.013*
I feel comfortable providing counseling to patients and/or families about medications for treatment of opioid use disorder including long-term monitoring and risk/benefits	13 (24.1%)	15 (46.9%)	0.006*

<b>Didactic Knowledge Scores Pre- and Post-Intervention Using McNemar Test</b>	<b>Pre-Intervention Correct, n(%)</b>	<b>Post-Intervention Correct, n(%)</b>	<b>p-value</b>
Which of the following is the correct scoring device for opioid withdrawal symptoms?	40 (72.7%)	31 (96.9%)	0.021*
The opioid withdrawal scoring tool includes which of the following?	29 (53.7%)	22 (68.6%)	0.289
Which of the following is the best treatment for a 16-year-old with opioid use disorder?	35 (66%)	30 (100%)	<0.001*
True or false: In Kentucky, confidentiality laws enforce disclosure of treatment for substance use disorder to a guardian.	38 (69.1%)	23 (71.9%)	1