

Case-Based Likert Confidence Scores Pre- and Post-Intervention Using Wicoxon Signed-Rank Test	Pre-Intervention Confident, n(%)	Post-Intervention Confident, n(%)	p-value
I can effectively screen for substance use disorder, including alcohol and opioids	9 (19.1%)	17 (47.2%)	0.091
I can recognize the symptoms of opioid withdrawal	11 (23.4%)	20 (55.6%)	0.03*
I can choose the most appropriate medication to treat opioid use disorder using my patient's age	13 (27.7%)	19 (52.8%)	0.001*
I can effectively communicate with families of my patients while maintaining their confidentiality	17 (36.2%)	16 (44.4%)	0.041*
I feel comfortable providing counseling to patients and/or families about medications for treatment of opioid use disorder including long-term monitoring and risk/benefits	15 (31.9%)	16 (44.4%)	0.156

Case-Based Knowledge Scores Pre- and Post Intervention Using McNemar Test	Pre-Intervention Correct, n(%)	Post-Intervention Correct, n(%)	p-value
Which of the following is the correct scoring device for opioid withdrawal symptoms?	30 (63.8%)	35 (100%)	<0.001*
The opioid withdrawal scoring tool includes which of the following?	24 (51.1%)	25 (71.4%)	0.146
Which of the following is the best treatment for a 16-year-old with opioid use disorder?	32 (68.1%)	24 (68.6%)	1
True or false: In Kentucky, confidentiality laws enforce disclosure of treatment for substance use disorder to a guardian	37 (78.7%)	24 (68.6%)	0.25