

Table 1. Demographic data for infants receiving autologous UCBC infusion, and contemporaneous cohort not receiving UCBC infusion. Data are expressed as median (IQR) or n (%).

Demographics	Infused infants N=23	Contemporaneous cohort N = 93
Perinatal		
Maternal age (years)	35 (30-37)	32 (29-35)
Maternal Body Mass Index	28 (23-32)	27 (24-33)
Pregnancy induced hypertension	4/23 (17.3)	11/93 (11.8)
Gestational diabetes	7/23 (30.4)	11/93 (11.8)
Chorioamnionitis	6/23 (26.0)	6/93 (6.4)
Prolonged rupture of membranes >18 hours	12/23 (52.1)	35/93 (37.6)
Antepartum haemorrhage	3/23 (13.0)	27/93 (29.0)
Fetal growth restriction	8/23 (34.7)	16/93 (17.2)
Antenatal steroids (complete course)	22/23 (95.6)	60/93 (64.5)
Magnesium sulphate at delivery (any)	23/23 (100)	79/93 (84.9)
Caesarean section	19/23 (82.6)	51/93 (54.8)
Deferred cord clamping, at least 60 seconds	16/23 (69.5)	66/93 (70.9)
Intubation at birth	6/23 (26.0)	27/93 (29.0)
Surfactant at birth	5/23 (21.7)	15/93 (16.1)
Neonatal		
Gestational age at birth (completed weeks)	26 (25-27)	26 (24-27)
Birth weight (grams)	750 (650-946)	769 (660-1017)
Male sex	17/23 (73.9)	39/93 (41.9)
Apgar score at 5 min	8 (6-9)	8 (7, 8)
IVH grade 1-2 (pre-infusion)	13/22 (59.0)	48/93 (51.6)
Patent ductus arteriosus treated (pre-infusion)	13/22 (59.0)	42/93 (45.1)