

**Table 2.** Post-Acute Sequelae of COVID-19 Index scores categorized based on proposed cut-points compared between compared based on index emergency department visit SARS-CoV-2 test result (i.e., positive vs. negative).

Age categories	Index	Total	SARS-CoV-2 Positive	SARS-CoV-2 Negative	Odds ratio (95%CI)*
<b>6-Month Follow-Up</b>					
Aged 6 to <12 years	$\geq 5.5$	60/258 (23.3)	29/115 (25.2)	31/143 (21.7)	1.22 (0.61, 2.44)
Aged 12 to <18 years	$\geq 5.0$	10/79 (12.7)	5/28 (17.9)	5/51 (9.8)	2.00 (0.49, 8.22)
<b>12-Month Follow-Up</b>					
Aged 6 to <12 years	$\geq 5.5$	144/479 (30.1)	41/152 (27.0)	103/327 (31.5)	0.83 (0.46, 1.40)
Aged 12 to <18 years	$\geq 5.0$	33/226 (14.6)	9/52 (17.3)	24/174 (13.8)	1.31 (0.49, 3.50)

\* Odds ratios were obtained from logistic regression fitted in generalized estimating equation (GEE) model with binary outcome being the PASC Index score exceeding the threshold level, with the model adjusted for clustering by site. Separate GEE models were used for each subgroup included in this table.