

Characteristic		OO/SO Group N=50 n (%)	SO Group N=51 n (%)	Total N=101 n (%)
Sex	Female	16 (32.0)	29 (56.9)	45 (44.6)
	Male	34 (68.0)	22 (43.1)	56 (55.4)
Race	White	19 (38.0)	23 (45.1)	42 (41.6)
Ethnic Group	Hispanic or Latino	3 (6.0)	5 (9.8)	8 (7.9)
	Not Hispanic or Latino	44 (88.0)	42 (82.4)	86 (85.1)
	Not Reported	3 (6.0)	4 (7.8)	7 (6.9)
Diagnosis for PN	Prematurity	46 (90.2)	46 (85.2)	92 (87.6)
	Gastroschisis	0	3 (5.6)	3 (2.9)
	Enteropathies including inflammatory gastrointestinal diseases	0	1 (1.9)	1 (1.0)
	Feeding intolerance	1 (2.0)	0	1 (1.0)
	Meconium ileus	1 (2.0)	0	1 (1.0)
	Necrotizing enterocolitis	1 (2.0)	0	1 (1.0)
	Persistent pulmonary hypertension	1 (2.0)	0	1 (1.0)
	Preterm infant with feeding intolerance	1 (2.0)	0	1 (1.0)
	Preterm infant with gastroschisis	0	1 (1.9)	1 (1.0)
	Sepsis	0	1 (1.9)	1 (1.0)
	Short bowel syndrome	0	1 (1.9)	1 (1.0)
	Small for gestational age	0	1 (1.9)	1 (1.0)
Age Group	Preterm Neonate (born <37 weeks of gestation)	47 (94.0)	47 (92.2)	94 (93.1)
	Full Term Neonate (born ≥37 weeks of gestation to <1 month of age)	2 (4.0)	2 (3.9)	4 (4.0)
	Infant (1 to <12 months of age)	0	0	0
	Child (1 to <10 years of age)	1 (2.0)	1 (2.0)	2 (2.0)
	Adolescent (10 to <18 years of age)	0	1 (2.0)	1 (1.0)
Weight at Birth	Normal (>2500g)	1 (2.0)	2 (3.9)	3 (3.0)
	Low (1500–2499g)	4 (8.0)	6 (11.8)	10 (9.9)
	Very Low (1000–1499g)	25 (50.0)	28 (54.9)	53 (52.5)
	Extremely Low (<1000g)	19 (38.0)	13 (25.5)	32 (31.7)
	N/A ¹	1 (2.0)	2 (3.9)	3 (3.0)

