

Table 1: Baseline and Clinical Characteristics of the Study Population

	Donor Milk (n=85)	Preterm Formula (n=83)
Baseline Characteristics		
Gestational age (weeks), median (IQR)	26 (25–27)	26 (25–27)
Birth weight (grams), median (IQR)	820 (630–980)	815 (650–939)
Male sex, n (%)	35 (41.2)	41 (49.5)
Small for gestational age, n (%)	12 (14.1)	8 (9.6)
Race/ethnicity		
White, n (%)	27 (31.8)	39 (47.0)
Black, n (%)	58 (68.2)	44 (53.0)
Hispanic, n (%)	0	0
Asian, n (%)	0	0
Other, n (%)	0	0
Multiple gestation, n (%)	21 (24.7)	26/83 (31.3)
Antenatal corticosteroids, n (%)	79 (92.9)	73/83 (88.0)
Cesarean delivery, n (%)	60 (70.5)	60 (72.3)
Surfactant exposure, n (%)	65 (76.5)	60 (72.3)
Clinical Characteristics		
Spontaneous intestinal perforation, n (%)	4 (4.7)	0
Bronchopulmonary dysplasia, n (%)	27 (35.1)	36 (46.2)
Patent ductus arteriosus, n (%)	26 (30.6)	31 (37.3)
Pharmacologically treated patent ductus arteriosus, n (%)	8 (9.4)	11 (13.3)
Intracranial hemorrhage (ICH), n (%)	21 (24.7)	20 (24.1)
Severe ICH, n (%)	8 (9.4)	11 (13.3)
Early onset sepsis, n (%)	2 (2.4)	4 (4.8)
Late onset sepsis, n (%)	19 (22.4)	11 (13.3)
Necrotizing enterocolitis, n (%)	3 (3.5)	8 (9.6)
Death, n (%)	7 (8.2)	5 (6.0)