

Table 2. Clinical Features of Cases and Controls (n=108).

	Case (n=50)	Controls (n=58)	p - value
DGBI before AGE (%)	4 (8%)	1 (1.7%)	0.179
DGBI Prevalence at 3 months (%)	10 (20%)	1 (1.7%)	0.002
IBS* (%)	7 (14%)	1 (1.7%)	
FD** (%)	1 (2%)	0	
Overlap IBS + FD (%)	2 (4%)	0	
Incident cases of DGBI at 3 months (%)	6 (12%)	0	0.008
IBS (%)	3 (6%)	0	
FD (%)	1 (2%)	0	
Overlap IBS + FD (%)	2 (4%)	0	
DGBI Prevalence at 6 months*** (%)	9 (18%)	1 (1.7%)	0.006
IBS (%)	7 (14%)	1 (1.7%)	
FD (%)	1 (2%)	0	
Overlap IBS + FD (%)	1 (2%)	0	
Incident cases of DGBI at 6 months*** (%)	5 (10%)	0	0.021
IBS (%)	3 (6%)	0	
FD (%)	1 (2%)	0	
Overlap IBS + FD (%)	1 (2%)	0	

* Irritable bowel syndrome (IBS)

** Functional dyspepsia (FD)

*** One was lost to follow-up