

**Table 1. Patient characteristics, stratified by hypertension status**

	2.1. By Hypertension (Yes/No)		2.1. By Hypertension stage				p-value
	No HTN (Normal + Elevated) (n=42)	HTN (Stage 1-2) (n=94)	Normal (n=12)	Elevated (n=30)	Stage 1 (n=54)	Stage 2 (n=40)	
<b>Sleep duration</b>							
Adequate	30 (71.4)	66 (70.2)	8 (66.7)	22 (73.3)	38 (70.4)	28 (70.0)	0.97
Short	12 (28.6)	28 (29.8)	4 (33.3)	8 (26.7)	16 (29.6)	12 (30.0)	
Age, median (IQR)	14 (11, 15)	14 (11, 16)	13 (10.5, 16)	14 (12, 15)	13 (10, 17)	15 (13, 16)	0.78
<b>Gender</b>							
Male	27 (64.3)	58 (61.7)	6 (50.0)	21 (70.0)	33 (61.1)	25 (62.5)	0.66
Female	15 (35.7)	36 (38.3)	6 (50.0)	9 (30.0)	21 (38.9)	15 (37.5)	
<b>Race</b>							
White	28 (66.7)	60 (63.8)	9 (75.0)	19 (63.3)	37 (68.5)	23 (57.5)	0.43
Black	10 (23.8)	22 (23.4)	1 (8.3)	9 (30.0)	9 (16.7)	13 (32.5)	
Asian	3 (7.1)	5 (5.3)	2 (16.7)	1 (3.3)	3 (5.6)	2 (5.0)	
Unknown	1 (2.4)	7 (7.4)	0 (0.0)	1 (3.3)	5 (9.3)	2 (5.0)	
<b>Ethnicity</b>							
Hispanic	22 (52.4)	43 (45.7)	5 (41.7)	17 (56.7)	27 (50.0)	16 (40.0)	0.82
Non Hispanic	19 (45.2)	45 (47.9)	7 (58.3)	12 (40.0)	24 (44.4)	21 (52.5)	
Unknown	1 (2.4)	6 (6.4)	0 (0.0)	1 (3.3)	3 (5.6)	3 (7.5)	
National ADI, median (IQR)	53.5 (31.0, 69.0)	57.0 (40.0, 71.0)	32.5 (22.0, 49.0)	59.5 (39.0, 74.0)	60.5 (40.0, 72.0)	55.0 (39.5, 62.5)	<b>0.03</b>
<b>ADI quartiles</b>							
Q1	8 (19.0)	13 (13.8)	4 (33.3)	4 (13.3)	7 (13.0)	6 (15.0)	0.36
Q2	11 (26.2)	19 (20.2)	5 (41.7)	6 (20.0)	9 (16.7)	10 (25.0)	
Q3	15 (35.7)	46 (48.9)	2 (16.7)	13 (43.3)	29 (53.7)	17 (42.5)	
Q4	8 (19.0)	16 (17.0)	1 (8.3)	7 (23.3)	9 (16.7)	7 (17.5)	