

Table 3. Adjusted Odds Ratios of FI Severity and Asthma Outcomes Stratified by Food Benefit Use
 [Weighted N=35,536,713 (3,638 Observations)]

| | Families with Benefits | | Families Without Benefits | |
|---|------------------------|--------------|---------------------------|--------------|
| | aOR [†] | 95% CI | aOR [†] | 95% CI |
| 3A. Outcome: Asthma Exacerbation | | | | |
| Food Insecurity (ref = High Food Security) | | | | |
| Marginal Food Security | 1.33 | [0.82,2.16] | 1.35 | [0.87, 2.10] |
| Low Food Security | 2.54*** | [1.64, 3.95] | 1.17 | [0.74,1.87] |
| Very Low Food Security | 3.50*** | [2.11,5.79] | 3.29** | [1.26, 4.19] |
| 3B. Outcome: ED/UC Visit for Asthma | | | | |
| Food Insecurity (ref = High Food Security) | | | | |
| Marginal Food Security | 1.17 | [0.63, 2.18] | 0.92 | [0.48, 1.77] |
| Low Food Security | 1.44 | [0.79, 2.64] | 0.99 | [0.50, 1.95] |
| Very Low Food Security | 4.27*** | [2.43, 7.47] | 2.27 | [0.93, 5.51] |

* p<0.05, ** p<0.01, *** p<0.001

[†]aOR: Adjusted for age, sex, race/ethnicity, insurance status, self-reported household income (>1.99% FPL or <1.99% FPL), self-reported parental education, and household structure.