

	<b>Overall,</b> N = 509 n (%)	<b>Facility Deaths</b> N = 364 n (%)	<b>Community Deaths</b> N = 145 n (%)	<i>P</i> value <sup>a</sup>
<b>Sex</b>				0.168
Male	279 (54.8)	207 (56.9)	72 (49.7)	
Female	230 (45.2)	157 (43.1)	73 (50.3)	
<b>Age at death</b>				0.654
Infants (aged 28-364 days)	287 (56.4)	208 (57.1)	79 (54.5)	
Young children (aged 12-59 months)	222 (43.6)	156 (42.9)	66 (45.5)	
<b>Age in days, median [IQR]</b>	294 [97, 631]	280 [90, 597]	314 [121, 726]	0.060
<b>Weight-for-age Z-score (N = 177)</b>				0.255
Normal ( $\geq$ -2SD)	159 (31.9)	113 (31.6)	46 (32.9)	
Moderate underweight (<-2 SD, -3 SD)	58 (11.6)	47 (13.1)	11 (7.9)	
Severe underweight (<-3 SD)	281 (56.4)	198 (55.3)	83 (59.3)	
<b>Length-for-age Z-score (N = 175)</b>				0.473
Normal ( $\geq$ -2SD)	238 (48.3)	164 (46.6)	74 (52.5)	
Moderate stunting (<-2 SD, -3 SD)	76 (15.4)	55 (15.6)	21 (14.9)	
Severe stunting (<- 3 SD)	179 (36.3)	133 (37.8)	46 (32.6)	
<b>Weight-for-length Z-score (N = 162)</b>				0.168
Normal ( $\geq$ -2SD)	190 (41.7)	143 (44.3)	47 (35.3)	
Moderate wasting (<-2 SD, -3 SD)	72 (15.8)	51 (15.8)	21 (15.8)	
Severe wasting (<-3 SD)	194 (42.5)	129 (39.9)	65 (48.9)	
<b>Site</b>				<0.001
Bangladesh	5 (1.0)	5 (1.4)	0 (0.0)	
Ethiopia	29 (5.7)	10 (2.7)	19 (13.1)	
Kenya	81 (15.9)	33 (9.1)	48 (33.1)	
Mali	14 (2.8)	7 (1.9)	7 (4.8)	
Mozambique	111 (21.8)	88 (24.2)	23 (15.9)	
Sierra Leone	108 (21.2)	102 (28.0)	6 (4.1)	
South Africa	161 (31.6)	119 (32.7)	42 (29.0)	