

**Table: HSHS St. John's NICU Very Preterm Infant Data (2007 – 2024; n = 1618 patients)**

<u>GA</u>	<u>#</u>	<u>Survived</u>	<u>BPD</u>	<u>LIVH, IPH, or cPVL</u>	<u>NEC</u>	<u>SIP</u>	<u>ROP</u>
23 – 31	1618	91%	24%	7% / 5%*	4%	1%	4%
23	106	49%	87%	20% / 12%*	4%	7%	14%
24	87	71%	81%	22% / 18%*	2%	9%	17%
25	125	80%	67%	16% / 13%*	7%	1%	13%
26	149	90%	50%	10% / 10%*	7%	1%	3%
27	158	94%	44%	8% / 7%*	5%	1%	3%
28	172	98%	19%	3% / 3%*	5%	none	1%
29	185	98%	10%	4% / 4%*	3%	none	2%
30	266	98%	3%	4% / 3%*	<1%	none	<1%
31	370	99%	1%	1% / 1%*	2%	none	<1%

Survived: alive at the time of NICU discharge

BPD: bronchopulmonary dysplasia requiring CPAP or oxygen at 36 weeks PMA

LIVH: large intraventricular hemorrhage with blood occupying  $\geq 50\%$  of the ventricular volume

cPVL: cystic periventricular leukomalacia

IPH: intraparenchymal hemorrhage

NEC:  $\geq$  stage II necrotizing enterocolitis

SIP: spontaneous intestinal perforation

ROP: retinopathy of prematurity requiring bevacizumab or laser surgery

\*Percent of survivors with LIVH, IPH, or cPVL for each row