

Table 1: Sociodemographic characteristics of patients by food insecurity (FI) at the congenital heart clinic

Characteristic	Total –includes all visit types (N= 4862)	Food Insecurity Positive (n=112)	Food Insecurity Negative (n=4750)	p-value
Age, mean +/-SD (years)	17.4 +/- 14.4	18.4 +/- 14.5	17.3 +/- 14.3	0.46
Male sex	2052(42.2%)	41 (36.6%)	1213 (40.3%)	0.44
Race				—
White	4,011 (82.5%)	79 (70.5%)	2,505 (83.1%)	
Black/African Am.	499 (10.3%)	20 (17.9%)	281 (9.3%)	
Asian	73 (1.5%)	1 (0.9%)	48 (1.6%)	
Other/Unknown	279 (5.7%)	12 (10.7%)	179 (5.9%)	
Hispanic ethnicity	268 (5.5%)	12 (10.7%)	170 (5.6%)	0.024
Preferred language: English	4,769 (98.1%)	107 (95.5%)	2,948 (97.8%)	0.11
Abbreviations: SD, Standard Deviation				
<p>Notes: Values are n (%) unless otherwise indicated. P-values from Pearson χ^2 tests or t-test. Hispanic vs non-Hispanic patients were more likely to report FI (OR=2.0, 95% CI:1.08–3.73), while Black or African American vs White or Caucasian patients were also more likely to report FI (OR=2.26, 95% CI:1.36–3.74).</p>				